

Spectral gap estimates for some block matrices

Ivan Veselić*, Krešimir Veselić †

Abstract

We estimate the size of the spectral gap at zero for some Hermitian block matrices. Included are quasi-definite matrices, quasi-semidefinite matrices (the closure of the set of the quasi-definite matrices) and some related block matrices which need not belong to either of these classes. Matrices of such structure arise in quantum models of possibly disordered systems with supersymmetry or graphene like symmetry. Some of the results immediately extend to infinite dimension.

1 Introduction

Consider (finite) Hermitian block matrices of the form

$$H = \begin{bmatrix} A & B \\ B^* & -C \end{bmatrix} \quad (1)$$

(the minus sign is set by convenience). If A, C are positive definite then the matrix H is called quasi-definite. These matrices have several remarkable properties, one of them is that they are always non singular with the spectral gap at zero

$$\rho(H) \supseteq (-\min \sigma(C), \min \sigma(A)) \quad (2)$$

(ρ the resolvent set, σ the spectrum). That is, the spectral gap of the block-diagonal part of H in (1) can only grow if any B is added. Moreover, quasi-definite matrices have two remarkable *monotonicity properties*

- (A) if B is replaced by tB , $t > 0$, then all the eigenvalues go monotonically asunder as t is growing ([13],[8]).
- (B) The same holds if A, C is replaced by $A + tI, C + tI$, $t > 0$, respectively ([6]).

In this note we study some related classes of matrices. If in (1) the blocks A, C are allowed to be only positive semidefinite then H will naturally be called *quasi-semidefinite*. These matrices need not be invertible.

It is relatively easy to characterise the nonsingularity of a quasi-semidefinite matrix, see Proposition 2.1 below. Giving estimates for the gap at zero is more involved and this note offers some results in this direction. Since the size of the spectral gap at zero is bounded from below by $2\|H^{-1}\|^{-1}$ we will give various bounds for this quantity in terms of the blocks A, B, C where A, C are only positive semidefinite and the properties of B come into play.

*TU-Chemnitz, Fakultät für Mathematik, 09107 Chemnitz, Germany,
e-mail: ivan.veselic@mathematik.tu-chemnitz.de

†Fernuniversität Hagen, Fakultät für Mathematik und Informatik Postfach 940, D-58084 Hagen, Germany, e-mail: kresimir.veselic@fernuni-hagen.de

It is known from [14] that the invertibility of B carries over to H but there no bound for H^{-1} was provided. Some bounds for H^{-1} were given in [15].

As a technical tool we derive a bound for the matrix $(I + AC)^{-1}$ with A, C positive semidefinite which may have independent interest. We also sketch a related functional calculus for such products. We then conjecture another bound for this matrix which is based on some special cases as well as on numerical evidence. More specifically, the present article provides the following.

1. A rigorous and a conjectured bound for $(I + AC)^{-1}$ with A, C positive semidefinite.
2. A characterisation of the nonsingularity of a quasi-semidefinite matrix.
3. A bound for H^{-1} based on the bound for B^{-1} including an immediate generalisation to unbounded Hilbert-space operators defined by quadratic forms. To this general environment we also extend an elegant estimate obtained by [9] for the special case $A = C$, $B = B^*$.
4. A bound for H^{-1} based on the geometry of the null-spaces of all of A, B, C and certain restrictions of these operators to the orthogonal complements of these null-spaces.
5. Several counterexamples; some of them showing that some plausibly looking generalisations of the properties (A), (B) above are not valid.
6. A study of the spectral gap of a particular class of matrices which arise in the quantum mechanical modelling of disordered systems (see e.g. [2]) where in (1) we have $C = A$ but A is not necessarily positive definite. In particular, we will illustrate how changing boundary conditions can remove spurious eigenvalues from the gap. This is a specific, thoroughly worked out example on how to deal successfully with what is called spectral pollution.

Let us remark, that the variety of special cases as well as techniques which we use, illustrate the fact that we did not succeed in obtaining a unified framework for spectral gap estimates for general quasi-definite matrices.

Quasi-semidefinite matrices and their infinite dimensional analogs have important applications in Mathematical Physics. While we have no space here to discuss the relevant models in detail, we would like to convey an impression of the questions arising in this context. They have been the motivation for much of the research presented here. Certain types of Dirac operators are important examples. In these cases the non-singularity of H is typically due to the one of B (see [15] where this phenomenon was dubbed 'off-diagonal dominance').

Another particular motivation are quantum mechanical models of disordered solids.

While this is a well established research field, recently there has been interest in such models which give rise to operators with block-structure, see e.g. [9] or [2]. For some of these models the block structure is a consequence of the Dirac-like symmetry arising in Hamiltonians describing graphene.

Let us describe some of the specific spectral features which are of interest in this context: We consider several instances of *one-parameter Hermitian pencils* $A + tB, t \in \mathbb{R}$. The well known *monotonicity property*, namely that the eigenvalues of $A + tB$ grow monotonically in t , if B is positive semidefinite can, at least partly, be carried over to quasi-definite matrices as show the properties (A), (B), listed above. Here a question of particular importance is whether and how fast the spectral gap increases as t grows. Several theorems of this paper provide answers to this question in specific situations.

As we mentioned, certain physical models of disordered systems give rise to block-structured operator families. In this context, estimates have to take into account two important aspects:

(I) The size of the original physical system is macroscopic, i.e. essentially infinite. A mathematical understanding of the physical situation is — as a rule — only possible to achieve by analysing larger and larger finite sample systems, that describe in the thermodynamic limit the original physical situation.

This leads to finite matrices or to operators with compact resolvent. In any case, effectively one can reduce the focus on a finite number, say n , of eigenvalues, when analysing monotonicity properties. However, n is not fixed but growing unboundedly, as one passes to larger and larger sample scales.

Thus, efficient estimates on spectral gaps (or derivatives of eigenvalues) are not allowed to depend on the system size — expressed in the dimension of the matrix or the number of eigenvalues n . We will pay special attention to this issue in the following.

(II) Due to the fact that one wants to model a disordered system, with a large number of degrees of freedom, there is in fact not just *one* coupling constant $\in \mathbb{R}$, but rather a whole collection of them $(\omega_j)_j$. Thus the considered operator pencil is originally of the form

$$A + \sum_j \omega_j B_j .$$

A one-parameter family arises if one freezes all coupling constants except one. As a consequence, one is not dealing with one fixed unperturbed operator A , but rather with whole collection of them, depending on the background configuration of the (other) coupling constants $(\omega_j, j \neq 0)$. For this reason it would be desirable to obtain estimates on the spectral gap which do not depend on specific features of A .

The plan of the paper is as follows: In the next section we provide certain basic preliminary estimates for quasi semi-definite matrices. In Section 3 the main results concerning the spectral gap size of such matrices are stated. These results are formulated for finite matrices. In Section 4 we explain which results carry immediately over to the setting of (possibly unbounded) operators defined as quadratic forms. This includes the mentioned generalisation of an estimate from [9], as well as a comparison with bounds obtained in [15]. Section 5 discusses a conjectured bound of the matrix $\|(I + AC)^{-1}\|$ which appears as a tool in the proof of Theorem 3.3. The last section considers a special class of finite difference matrices, not necessarily quasi-definite, studied in [2]. Here we show that a stable spectral gap at zero can be achieved by an appropriate tuning of boundary conditions. Similar phenomena, yet without rigorous proofs, are numerically observed on related models with random diagonal entries.

2 Some preliminary results

To set the stage we collect some rather elementary statements and estimates.

Proposition 2.1 *A quasi-semidefinite matrix*

$$H = \begin{bmatrix} A & B \\ B^* & -C \end{bmatrix} .$$

is singular, if and only if at least one of the subspaces

$$\mathcal{N}(A) \cap \mathcal{N}(B^*), \quad \mathcal{N}(C) \cap \mathcal{N}(B)$$

(\mathcal{N} denoting the null-space) is non-trivial. Moreover, in the obvious notation,

$$\mathcal{N}(H) = \begin{bmatrix} \mathcal{N}(A) \cap \mathcal{N}(B^*) \\ \mathcal{N}(C) \cap \mathcal{N}(B) \end{bmatrix}. \quad (3)$$

The value $\min(\sigma(A))$ is an eigenvalue of H if and only if $\mathcal{N}(B^*) = \{0\}$ (and similarly for $\min(-\sigma(C))$).

Proof. The equations

$$Ax + By = 0, \quad B^*x - Cy = 0$$

imply

$$x^*Ax + x^*By = 0, \quad y^*B^*x - y^*Cy = 0.$$

Since both x^*Ax and y^*Cy are real and non-negative, the same is true of $\pm x^*By$ such that, in fact, all three expressions vanish. Since A, C are Hermitian positive semidefinite this implies $Ax = 0$ and $Cy = 0$, then also $B^*x = 0$ and $By = 0$. This proves (3); for the last assertion apply (3) to the matrix $H - \min(\sigma(A))I$. Other assertions follow trivially. Q.E.D.

Corollary 2.2 *The matrix H is non-singular if and only if the matrices*

$$A + \sqrt{BB^*}, \quad C + \sqrt{B^*B}$$

are positive definite.

Corollary 2.3 *The null-space of the matrix H from (1) does not change, if A is replaced by tA , $t > 0$ (and similarly with B, C).*

To quantify the influence of B in (1) on the spectral gap in the quasi-definite case we proceed as follows. First note the fundamental equality, valid for all selfadjoint operators

$$\|(H - \lambda I)^{-1}\| = \text{dist}(\lambda, \sigma(H)). \quad (4)$$

Taking any λ from the open interval $(-\min \sigma(C), \min \sigma(A))$ we have

$$H - \lambda I = \begin{bmatrix} (A - \lambda I)^{1/2} & 0 \\ 0 & (C + \lambda I)^{1/2} \end{bmatrix} W \begin{bmatrix} (A - \lambda I)^{1/2} & 0 \\ 0 & (C + \lambda I)^{1/2} \end{bmatrix}$$

with

$$W = \begin{bmatrix} I & Z \\ Z^* & -I \end{bmatrix}, \quad Z = (A - \lambda I)^{-1/2} B (C + \lambda I)^{-1/2}.$$

As it is readily seen the eigenvalues of the matrix W are $\pm \sqrt{1 + \sigma_i^2}$, where σ_i are the singular values of Z (cf. [11]). Thus,

$$\|W\| = \sqrt{\|I + Z^*Z\|}, \quad \|W^{-1}\| = \sqrt{\|(I + Z^*Z)^{-1}\|}$$

This gives the estimate

$$\|(H - \lambda I)^{-1}\| \leq \frac{\sqrt{\|(I + Z^*Z)^{-1}\|}}{\min\{\min(\sigma(A) - \lambda), \min(\sigma(C) + \lambda)\}}. \quad (5)$$

Therefore by taking e.g. $\lambda = \lambda_0 = \frac{1}{2}(\min(\sigma(A) - \min(\sigma(C)))$ we obtain

$$\|(H - \lambda_0 I)^{-1}\| \leq \frac{2\sqrt{\|(I + Z^*Z)^{-1}\|}}{\min(\sigma(A) + \min(\sigma(C))).} \quad (6)$$

We see that the gap is stretched at least by the factor

$$\|(I + Z^*Z)^{-1}\|^{-1/2} = \sqrt{1 + \min_i \sigma_i^2}.$$

3 Spectral bounds for quasi-semidefinite matrices.

We will be particularly interested in how the appearance of the block B can create spectral gap at zero, if A, C alone are unable to do so. The size of this gap is bounded from below by the quantity $2/\|H^{-1}\|$, cf. (4).

As a preparation we will consider the matrices of the form $I + AC$ with A, C positive semidefinite. They will play a key role in our estimates and may have an independent interest of their own. Note that they are generally not Hermitian.

Theorem 3.1 *Let A, C be Hermitian positive semidefinite. Then*

$$(i) \quad \sigma(AC) = \sigma(CA) \subseteq [0, \infty). \quad (7)$$

$$(ii) \quad \|(I + AC)^{-1}\| \leq 1 + \frac{\min\{\|A\|^{1/2}\|L^*C\|, \|C\|^{1/2}\|AM\|\}}{1 + \min \sigma(AC)} \quad (8)$$

where

$$A = LL^*, \quad C = MM^*. \quad (9)$$

(iii) *The matrix AC is diagonalisable.*

Proof. The statements (i), (iii) above are not new (see [4], [5]) respectively. To prove (ii) note

$$\begin{aligned} (\lambda + AC)^{-1} &= \frac{1}{\lambda}I - \frac{1}{\lambda}(\lambda I + AC)^{-1}AC = \frac{1}{\lambda}I - \frac{1}{\lambda}(\lambda I + LL^*C)^{-1}LL^*C \\ &= \frac{1}{\lambda}I - \frac{1}{\lambda}L(\lambda I + L^*CL)^{-1}L^*C, \end{aligned} \quad (10)$$

So the spectra of CA and L^*CL coincide — up to possibly the point zero. Now, L^*CL is Hermitian positive semidefinite, hence

$$\|(I + L^*CL)^{-1}\| = \frac{1}{1 + \min \sigma(L^*CL)} = \frac{1}{1 + \min \sigma(AC)}$$

and therefore

$$\|(I + AC)^{-1}\| \leq 1 + \frac{\|L\|\|L^*C\|}{1 + \min \sigma(AC)}.$$

The second half of (8) is similar. Q.E.D.

From (8) it immediately follows

$$\|(I + AC)^{-1}\| \leq 1 + \frac{\|A\|^{1/2}\|C\|^{1/2}\|L^*M\|}{1 + \min \sigma(AC)} \leq 1 + \|A\|\|C\| \quad (11)$$

Proposition 3.2 *Let A, C be Hermitian positive semidefinite. Then*

$$\|I + AC\| \geq 1 \quad (12)$$

and the equality is attained if and only if $AC = 0$.

Proof. Since the norm dominates the spectral radius, and by (7) the latter is not less than one, (12) follows. In the case of equality the whole spectrum consists of the single point 1 that is, the spectrum of $AC = A^{1/2}A^{1/2}C$ is $\{0\}$. Then the spectrum of the Hermitian matrix $A^{1/2}CA^{1/2}$ also equals $\{0\}$. Hence $A^{1/2}CA^{1/2} = (C^{1/2}A^{1/2})^*C^{1/2}A^{1/2} = 0$ and then also $AC = (CA)^* = 0$. Q.E.D.

Theorem 3.3 *Let in (1) the matrix B be square and invertible and let*

$$\alpha = \sup_{x \neq 0} \frac{x^* A x}{x^* \sqrt{B B^*} x}, \quad \gamma = \sup_{x \neq 0} \frac{x^* C x}{x^* \sqrt{B^* B} x}. \quad (13)$$

Then

$$\|H^{-1}\| \leq \|B^{-1}\| (1 + \max\{\alpha, \gamma\} + \alpha\gamma) \quad (14)$$

Proof. Using the polar decomposition $B = U\sqrt{B^*B} = \sqrt{B B^*}U$ the following factorisation holds

$$H = \begin{bmatrix} (B B^*)^{1/4} & 0 \\ 0 & (B^* B)^{1/4} \end{bmatrix} \begin{bmatrix} \tilde{A} & U \\ U^* & -\tilde{C} \end{bmatrix} \begin{bmatrix} (B B^*)^{1/4} & 0 \\ 0 & (B^* B)^{1/4} \end{bmatrix} \quad (15)$$

with

$$\tilde{A} = (B B^*)^{-1/4} A (B B^*)^{-1/4}, \quad \tilde{C} = (B^* B)^{-1/4} C (B^* B)^{-1/4}.$$

Also,

$$\begin{bmatrix} \tilde{A} & U \\ U^* & -\tilde{C} \end{bmatrix}^{-1} = \begin{bmatrix} (I + \tilde{C}\tilde{A})^{-1}\tilde{C} & U(I + \tilde{C}\tilde{A})^{-1} \\ (I + \tilde{A}\tilde{C})^{-1}U^* & -\tilde{A}(I + \tilde{C}\tilde{A})^{-1} \end{bmatrix} \quad (16)$$

where

$$\hat{A} = U^* \tilde{A} U, \quad \hat{C} = U \tilde{C} U^*$$

are again Hermitian positive semidefinite. This is immediately verified taking into account the identity

$$(I + \hat{A}\hat{C})^{-1}U^* = U^*(I + \tilde{A}\tilde{C})^{-1}. \quad (17)$$

This, together with the identities of the type

$$(I + \hat{C}\hat{A})^{-1}\hat{C} = \hat{C}^{1/2}(I + \hat{C}^{1/2}\hat{A}\hat{C}^{1/2})^{-1}\hat{C}^{1/2} \quad (18)$$

and the obvious inequality

$$\left\| \begin{bmatrix} E & F \\ G & K \end{bmatrix} \right\| \leq \max\{\|E\|, \|K\|\} + \max\{\|F\|, \|G\|\}$$

permits the use of (11) and the factorisation (15) to obtain (14). Here we have used obvious identities

$$\alpha = \|\tilde{A}\| = \|\hat{A}\|, \quad \gamma = \|\tilde{C}\| = \|\hat{C}\|$$

and the fact that U is unitary. Q.E.D.

If B is replaced by tB , $t > 0$ then (14) goes over into

$$\|H^{-1}\| \leq \frac{\|B^{-1}\|}{t} \left(1 + \frac{\max\{\alpha, \gamma\}}{t} + \frac{\alpha\gamma}{t^2} \right) \quad (19)$$

Note that here the right hand side is monotone decreasing in t . The larger t , the larger the gap at zero.

The proof of Theorem 3.3 may appear odd: the estimate for the inverse of a Hermitian matrix H relies heavily on the estimate for the inverse of some *non-Hermitian* matrices of the type $I + AC$. But this is the price for halving the dimension of the problem in working with 'non-symmetric' Schur complements.

On the other hand, if both A , C are positive definite then setting in (16) $\tilde{C} = \hat{C} = C$, $\tilde{A} = \hat{A} = A$, $U = I$, (2) yields

$$\|(I + AC)^{-1}\| \leq \max\{\|A^{-1}\|, \|C^{-1}\|\}. \quad (20)$$

Remark 3.4 By (7) the spectrum of $I + AC$ is uniformly bounded away from zero, so one may ask whether there is a uniform bound for the norm of its inverse. The answer is negative as shows the following example which is due to M. Omladič (private communication). Set

$$A = \begin{bmatrix} t & 0 \\ 0 & 1/t \end{bmatrix}, \quad C = \begin{bmatrix} 1/t & 1 \\ 1 & t \end{bmatrix}.$$

Then

$$I + AC = \begin{bmatrix} 2 & t \\ 1/t & 2 \end{bmatrix}, \quad (I + AC)^{-1} = \frac{1}{3} \begin{bmatrix} 2 & -t \\ -1/t & 2 \end{bmatrix}, \quad (21)$$

and this is not bounded as t varies over positive reals.

We now turn to a more complicated case in which A, C may have null-spaces and the invertibility of H is due to the conspiring of all three blocks A, B, C . As an additional information we will need lower bounds for the non-vanishing part of $\sigma(A), \sigma(C)$. Thus, it will be technically convenient to represent H in a block form which explicitly displays these null-spaces:

$$H = \begin{bmatrix} A & 0 & B_{11} & B_{12} \\ 0 & 0 & B_{21} & B_{22} \\ B_{11}^* & B_{21}^* & -C & 0 \\ B_{12}^* & B_{22}^* & 0 & 0 \end{bmatrix}. \quad (22)$$

Here, for the notational simplicity, the new blocks A, C are the 'positive definite restrictions' of the original blocks A, C in (1). In view of Proposition 2.1 H is non-singular, if and only if both matrices

$$\begin{bmatrix} B_{21} & B_{22} \end{bmatrix}, \quad \begin{bmatrix} B_{12} \\ B_{22} \end{bmatrix}$$

have full rank. The following theorem gives a new sufficient condition for invertibility and subsequently a gap estimate.

Theorem 3.5 *Suppose that*

$$H = \begin{bmatrix} A & B \\ B^* & -C \end{bmatrix}.$$

is quasi-semidefinite. Assume, in addition,

1. $\dim(\mathcal{N}(A)) = \dim(\mathcal{N}(C))$
2. B is a one-to-one map from $\mathcal{N}(C)$ onto $\mathcal{N}(A)$.

That is, the block B_{22} in (22) is square and non-singular. Then

$$(-\varepsilon, \varepsilon) \cap \sigma(H) = \emptyset$$

with

$$\varepsilon \geq \frac{1}{(1 + \max\{\|B_{12}B_{22}^{-1}\|, \|B_{21}^*B_{22}^{-*}\|\})^2 \max\{\|A^{-1}\|, \|C^{-1}\|, \|B_{22}^{-1}\|\}}.$$

Proof. We represent H in the unitarily equivalent, permuted form

$$\begin{bmatrix} A & B_{11} & 0 & B_{12} \\ B_{11}^* & -C & B_{21}^* & 0 \\ 0 & B_{21} & 0 & B_{22} \\ B_{12}^* & 0 & B_{22}^* & 0 \end{bmatrix} = \begin{bmatrix} \hat{A} & \hat{B} \\ \hat{B}^* & \hat{G} \end{bmatrix}. \quad (23)$$

By renaming this matrix again into H we now perform the decomposition

$$H = \begin{bmatrix} I & \widehat{B}\widehat{G}^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} \widehat{A} - \widehat{B}\widehat{G}^{-1}\widehat{B}^* & 0 \\ 0 & \widehat{G} \end{bmatrix} \begin{bmatrix} I & 0 \\ \widehat{G}^{-1}\widehat{B}^* & I \end{bmatrix}.$$

Hence the simple estimate

$$\|H^{-1}\| \leq (1 + \|\widehat{B}\widehat{G}^{-1}\|)^2 \max\{\|(\widehat{A} - \widehat{B}\widehat{G}^{-1}\widehat{B}^*)^{-1}\|, \|\widehat{G}^{-1}\|\}.$$

We now bound the single factors above:

$$\begin{aligned} \widehat{A} - \widehat{B}\widehat{G}^{-1}\widehat{B}^* &= \widehat{A} - \begin{bmatrix} 0 & B_{12} \\ B_{21}^* & 0 \end{bmatrix} \begin{bmatrix} 0 & B_{22}^{-*} \\ B_{22}^{-1} & 0 \end{bmatrix} \begin{bmatrix} 0 & B_{21} \\ B_{12}^* & 0 \end{bmatrix} \\ &= \begin{bmatrix} A & B_{11} - B_{12}B_{22}^{-1}B_{21} \\ B_{11}^* - B_{21}^*B_{22}^{-*}B_{12}^* & -C \end{bmatrix}. \end{aligned}$$

This matrix is quasi-definite, hence the interval $(-\min \sigma(C), \min \sigma(A))$ contains none of its eigenvalues. Thus, $\widehat{A} - \widehat{B}\widehat{G}^{-1}\widehat{B}^*$ is invertible and

$$\|(\widehat{A} - \widehat{B}\widehat{G}^{-1}\widehat{B}^*)^{-1}\| \leq \max\{\|A^{-1}\|, \|C^{-1}\|\}.$$

Furthermore,

$$\|\widehat{G}^{-1}\| = \|B_{22}^{-1}\|$$

and

$$\widehat{B}\widehat{G}^{-1} = \begin{bmatrix} B_{12}B_{22}^{-1} & 0 \\ 0 & B_{21}^*B_{22}^{-*} \end{bmatrix},$$

hence

$$\|H^{-1}\| \leq (1 + \max\{\|B_{12}B_{22}^{-1}\|, \|B_{21}^*B_{22}^{-*}\|\})^2 \max\{\|A^{-1}\|, \|C^{-1}\|, \|B_{22}^{-1}\|\}.$$

Q.E.D.

Note that the radius of the resolvent interval, guaranteed in the the previous theorem depends on the spectra of some operators, obtained from the original blocks A, B, C .

If in the preceding theorem we replace B by tB and t is sufficiently large then we obtain

$$\varepsilon \geq \frac{t}{(1 + \max\{\|B_{12}B_{22}^{-1}\|, \|B_{21}^*B_{22}^{-*}\|\})^2 \|B_{22}^{-1}\|}. \quad (24)$$

Another relevant special case has $A = C$ and $B^* = B$, both positive definite. Then, as was shown in [9], we have

$$\rho(H) \supseteq (-\sqrt{\min \sigma(A)^2 + \min \sigma(B)^2}, \sqrt{\min \sigma(A)^2 + \min \sigma(B)^2}). \quad (25)$$

Remark 3.6 The technique used in the proof of Theorem 3.1 is related to the more general functional calculus for products AC , with A, C bounded, selfadjoint, C positive semidefinite in a general Hilbert space. It reads

$$f \mapsto f(AC) = f(0)I + AC^{1/2}f_1(C^{1/2}AC^{1/2})C^{1/2} \quad (26)$$

with

$$f_1(\lambda) = \begin{cases} \frac{f(\lambda) - f(0)}{\lambda}, & \lambda \neq 0 \\ f'(0), & \lambda = 0 \end{cases}$$

By the property

$$f(XY)X = Xf(YX), \quad (27)$$

valid for any matrix analytic function f , this obviously extends the standard analytic functional calculus and requires f to be differentiable at zero, otherwise just bounded measurable; then f_1 will again be bounded measurable and applied to a selfadjoint operator $C^{1/2}AC^{1/2}$.¹ This calculus is a Hilbert-space generalisation of the assertions of Theorem 3.1, (i), (ii), only here the point zero may remain a sort of a 'spectral singularity'.

The linearity and multiplicativity of the map $f \mapsto f(AC)$ is verified by straightforward algebraic manipulations. Also, if the functions f in (26) are endowed with the norm

$$\|f\| = |f(0)| + \|f_1\|_\infty \quad (28)$$

then the map $f \mapsto f(AC)$ is obviously continuous. This admits estimating some other interesting functions of AC , for instance, the group e^{-ACt} , if both A and C are positive semidefinite and $t > 0$. In this case $f(\lambda) = e^{-t\lambda}$ and it is immediately verified that $|f_1|$, $\lambda \geq 0$, is bounded by t , hence

$$\|e^{-ACt}\| \leq 1 + t\|AC^{1/2}\|\|C^{1/2}\|. \quad (29)$$

and similarly

$$\|e^{-ACt}\| \leq 1 + t\|CA^{1/2}\|\|A^{1/2}\|. \quad (30)$$

Remark 3.7 Extending monotonicity properties? In the introduction we have stated two known monotonicity properties of the eigenvalues for some affine quasi-definite pencils. It is natural to try to extend this monotonicity to some neighbouring classes of matrix families. Some of our examples will be of the form

$$H = \begin{bmatrix} A & B \\ B^* & -A \end{bmatrix} \quad (31)$$

with 2×2 -matrices $A^* = A$ and $B^* = \pm B$ and no (semi)definiteness assumption whatsoever. Here a straightforward calculation shows that the characteristic polynomial is

$$\lambda^4 - 2(\|A\|_F^2 + \|B\|_F^2)\lambda^2 + |\det(A - \sqrt{\mp 1}B)|^2 \quad (32)$$

where $\|\cdot\|_F$ means the Frobenius or Hilbert-Schmidt norm. This can be used to give a general formula for the roots explicitly, see the Appendix.

If in a quasi-definite matrix (1) the matrices A and C increase (in the sense of forms), then the estimate (2) certainly improves, but does the gap at zero also necessarily increase? The answer is no as the following example due to W. Kirsch (private communication), shows. Set

$$H = \begin{bmatrix} A & B_t \\ B_t & -A \end{bmatrix} \quad (33)$$

with

$$A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}, \quad B_t = \begin{bmatrix} 1 & 0 \\ 0 & t \end{bmatrix}.$$

The matrix is quasi-definite. By (32) the characteristic equation is readily found to be

$$\lambda^4 - (11 + t^2)\lambda^2 + 13 + 5t^2 + 2t = 0 \quad (34)$$

¹This functional calculus, probably well-known by now, was described to the second author by the late C. Apostol, Bucharest, more than forty years ago.

and the absolutely smaller eigenvalue is given in Figure 1 as function of t , $5 < t < 20$, (conveniently scaled) and it shows a non-monotonic behaviour. Thus, there does not seem to be a simple generalisation of the monotonicity property (A). On the other hand by the unitary similarity

$$\begin{bmatrix} A & B_t \\ B_t & -A \end{bmatrix} = \frac{1}{2} \begin{bmatrix} I & I \\ I & -I \end{bmatrix} \begin{bmatrix} B_t & A \\ A & -B_t \end{bmatrix} \begin{bmatrix} I & I \\ I & -I \end{bmatrix} \quad (35)$$

the same holds for the property (B). Another likely generalisation of (B), namely to have




Figure 1: Lack of monotonicity

monotone eigenvalues, if in (31) the matrix A is replaced by tA , is also false. The counterexample is a numerical one:

$$A = \begin{bmatrix} 1.24 & 0.81 \\ 0.81 & 0.53 \end{bmatrix}, \quad B = \begin{bmatrix} 0.30 & -0.27 \\ -0.31 & -0.48 \end{bmatrix}.$$

Here both A and B are positive definite. The absolutely smaller eigenvalues for $5 < t < 20$ are shown in the Figure 2.

Finally, a yet simpler quasi-semidefinite example is given by

$$H = \begin{bmatrix} A_t & B \\ -B & -A_t \end{bmatrix} \quad (36)$$

with

$$A_t = \begin{bmatrix} 0 & 0 \\ 0 & t \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Note that $\det(H) = 1$ for all t and the spectrum $-\lambda_1, -\lambda_2, \lambda_2, \lambda_1$ is symmetric w.r.t. zero. Thus if $|\lambda_1|$ increases, $|\lambda_2|$ has to decrease with growing $t > 0$. This already shows that property (B) in the introduction cannot hold for $t \mapsto A + t \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. Moreover, it turns out that the spectral gap of (36) shrinks to zero as $t \rightarrow \infty$.




Figure 2: Lack of monotonicity

4 Unbounded operator matrices

Most of the results obtained above immediately extend to infinite dimensional Hilbert space. Theorem 3.1 (except (iii)) and Proposition 3.2, together with their proofs apply literally to any bounded selfadjoint positive semidefinite operators A, C . Theorem 3.3 even allows B, A, C as unbounded. In fact, the last two may be just quadratic forms, requiring, of course, that the quantities α, γ in (13), reformulated in the quadratic form context, be finite, whereas B has a bounded, everywhere defined inverse. More precisely, the operator H is defined by the *form block matrix*

$$\begin{bmatrix} \mathbf{a} & B \\ B^* & -\mathbf{c} \end{bmatrix} \quad (37)$$

where the symmetric sesquilinear forms \mathbf{a}, \mathbf{c} have to be defined on the form domains of $\sqrt{BB^*}, \sqrt{B^*B}$, respectively, and the relative form bounds

$$\alpha = \sup_{x \neq 0} \frac{\mathbf{a}(x, x)}{\|(B^*B)^{1/4}x\|^2}, \quad \gamma = \sup_{x \neq 0} \frac{\mathbf{c}(x, x)}{\|(BB^*)^{1/4}x\|^2}. \quad (38)$$

are finite. So, the operators $\tilde{A}, \tilde{C}, \hat{A}, \hat{C}$ appearing in the proof of Theorem 3.3 will again be bounded and positive semidefinite whereas the formula (15) now serves as a natural definition of the operator H itself. Indeed, H is given as a product of three selfadjoint operators, each having a bounded, selfadjoint inverse. The bounded invertibility of the first and the third factor in (15) is trivial, whereas for the second it follows from the formula (16) (cf. also [14]). Similar remarks hold for Theorem 3.5 as well. (Proposition 2.1 could also be reformulated in infinite dimension, but this will not interest us here.)

We will now compare our bound with a bound obtained in [15]. This bound (with our notation) requires that A, C be relatively bounded with respect to B, B^* , respectively. According to [7], Ch. VI, Th. 1.38 the operator boundedness implies the form boundedness with the same bound; so our setting is more general. Also [15] gives an eigenvalue bound under the condition that at least one of the operators A, C be bounded. The estimate obtained there is rather complicated; but if both A, C are bounded then [15] gives a somewhat

simpler bound

$$|\sigma(H)| \geq -\frac{1}{2}(\|A\| + \|C\|) + \left(\frac{1}{4}(\|A\| - \|C\|)^2 + (\|B^{-1}\|^{-2})\right)^{1/2} \quad (39)$$

which is still not easily compared with our estimate (14). Anyhow, if $\|A\| = \|C\|$ and the relative bound $\|A\|\|B^{-1}\|$ is larger than one, then the right hand side of (39) becomes negative and the estimate is void whereas the bound (14) always makes sense. In fact, our relative bounds α and γ may be arbitrary. In particular, they need not be less than one which is a usual requirement in operator perturbation theory. It is a general feature with quasi-definite matrices that perturbations, as long as they respect in an appropriate sense the block structure, need to be relatively bounded, but not necessarily with the relative bound less than one, in order to yield an effective perturbation theory. Such a phenomenon was already encountered, e. g. in [14].

The selfadjointness of the operator H from (37) immediately applies to various kinds of Dirac operators with supersymmetry (see [12], Sect. 5.4.2 and 5.5) under the appropriate definiteness assumption for the diagonal blocks.

An analogous construction of a selfadjoint block operator matrix H was made in [14] in the 'dual' case in which B is dominated by A, C in the sense that $A^{-1/2}BC^{-1/2}$ is bounded. Estimate (6) extends to this more general situation, where A, B, C need not be bounded.

Finally we come back to the estimate (25). The proof given in [9] went through squaring the matrix

$$H = \begin{bmatrix} A & B \\ B & -A \end{bmatrix} \quad (40)$$

which is inconvenient if A, B are unbounded. We provide an alternate proof under weaker assumption, namely that instead of operators A, B we have symmetric positive semidefinite (not necessarily closed) sesquilinear forms a, b defined on a dense domain $\mathcal{D} = \mathcal{D}(a) = \mathcal{D}(b)$. The obvious generalisation of the block operator matrix (40) is the symmetric sesquilinear form h , defined as

$$h(x, y) = a(x_1, y_1) + b(x_2, y_1) + b(x_1, y_2) - a(x_2, y_2), \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \quad (41)$$

for

$$x, y \in \mathcal{D} \oplus \mathcal{D}.$$

Neither of the forms a, b need be closed but *their sum* shall be assumed as closed. Here we have, in fact, first to construct the operator H . To this end we use the 'off-diagonalizing' transformation given by the unitary matrix

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ iI & -iI \end{bmatrix} \quad (42)$$

(cf. [12]). Obviously $U(\mathcal{D} \oplus \mathcal{D}) = \mathcal{D} \oplus \mathcal{D}$, whereas a direct calculation leads to

$$\begin{aligned} \widehat{h}(x, y) &= h(Ux, Uy) = ia(x_1, y_2) + b(x_1, y_2) + b(x_2, y_1) - ia(x_2, y_1) \\ &= -i\tau(x_2, y_1) + i\tau^*(x_1, y_2) \end{aligned} \quad (43)$$

where the forms

$$\tau = a - ib, \quad \tau^* = a + ib \quad (44)$$

are sectorial and mutually adjoint. Obviously the range of the form τ lies in the lower right quadrant of the complex plane.

The form τ is closed. This is readily seen from the equivalence of the corresponding norms:

$$\begin{aligned} |\tau(x, x)| &= \sqrt{a(x, x)^2 + b(x, x)^2} \leq a(x, x) + b(x, x) \leq \sqrt{2(a(x, x)^2 + b(x, x)^2)} \\ &= \sqrt{2}|\tau(x, x)| \end{aligned}$$

so that the closedness of $a + b$ is, in fact, equivalent to that of τ . Thus τ, τ^* generate mutually adjoint maximal sectorial operators T, T^* , respectively (see [7], Ch. VI, Theorem 2.1). Now, for $x_2 \in \mathcal{D}(T), x_1 \in \mathcal{D}(T^*)$ and $y_1, y_2 \in \mathcal{D}$ we have

$$\widehat{h}(x, y) = (\widehat{H}x, y) \quad (45)$$

where the operator

$$\widehat{H} = \begin{bmatrix} 0 & -iT \\ iT^* & 0 \end{bmatrix} \quad (46)$$

is obviously selfadjoint with the domain $\mathcal{D}(T^*) \oplus \mathcal{D}(T)$. Also selfadjoint is its inverse conjugate

$$H = U^* \widehat{H} U$$

with

$$h(x, y) = (Hx, y), \quad \mathcal{D}(H) \subseteq \mathcal{D}(\tau) \oplus \mathcal{D}(\tau). \quad (47)$$

The operator H is uniquely determined by (47) as is shown in [14], Proposition 2.3.

To estimate the inverse note that

$$\|Tz\| \|z\| \geq |(Tz, z)| = \sqrt{a(z, z)^2 + b(z, z)^2} \geq \sqrt{\alpha^2 + \beta^2} \|z\|^2$$

where $\alpha, \beta \geq 0$ is the lower bound of a, b , respectively. The above 'Lax-Milgram inequalities' are, in fact, the key argument in this matter. They are non-trivial, if any of α, β is different from zero. In this case by the maximality of T , its inverse is everywhere defined and

$$\|T^{-1}\| = \|T^{-*}\| \leq \frac{1}{\sqrt{\alpha^2 + \beta^2}}$$

By the obvious formula

$$\widehat{H}^{-1} = \begin{bmatrix} 0 & -iT^{-*} \\ iT^{-1} & 0 \end{bmatrix} \quad (48)$$

we finally obtain

$$\|\widehat{H}^{-1}\| = \|H^{-1}\| = \|T^{-1}\| \leq \frac{1}{\sqrt{\alpha^2 + \beta^2}} \quad (49)$$

which obviously reduces to (25) if a, b are bounded. Thus, we have proved the following theorem.

Theorem 4.1 *Let a, b be positive semidefinite symmetric sesquilinear forms with the common dense domain \mathcal{D} and respective lower bounds α, β and such that $a + b$ is closed. Then the form h from (41) defines a unique selfadjoint operator H with $\mathcal{D}(H) \subseteq \mathcal{D} \oplus \mathcal{D}$ and $h(x, y) = (Hx, y)$, for $x \in \mathcal{D}(H), y \in \mathcal{D} \oplus \mathcal{D}$. Moreover, if any of α, β is non-zero then H has a bounded inverse with*

$$\|H^{-1}\| \leq \frac{1}{\sqrt{\alpha^2 + \beta^2}}.$$

Remark 4.2 (i) The conditions of the preceding theorem are obviously fulfilled, if one of the forms a , b is closed and the other is relatively bounded with respect to the first. Moreover, if say, b is relatively bounded with respect to a then b need not be semidefinite; indeed the whole construction of H, \widehat{H}, T in the proof of the preceding theorem, goes through and we have

$$\|\widehat{H}^{-1}\| = \|H^{-1}\| = \|T^{-1}\| \leq \frac{1}{\alpha}, \quad (50)$$

provided that a is positive definite.

(ii) The form τ , constructed in the proof of the preceding theorem is not sectorial in the strict sense as defined in [7] because its range does not lie symmetrically with respect to the positive real axis. But, of course, the whole theory developed in [7] naturally and trivially extends to all kind of numerical ranges having semi-angle less than $\pi/2$. The standard form can be achieved simply by multiplying τ with a phase factor:

$$e^{\frac{\pi}{4}i}\tau = \frac{1}{\sqrt{2}}((a+b) + i(a-b)).$$

The symmetric part of this form is closed whereas the skew-symmetric part is relatively bounded with respect to the symmetric one, so it is sectorial in the strict sense of [7].

(iii) The obvious fact that the eigenvalues (whenever existing) of H are \pm singular values of T may have advantage in numerical computations with finite matrices. Firstly, the size of T is half the size of H and, secondly, there is a plenty of reliable computational software to compute the singular values (and vectors) of arbitrary matrices.

(iv) If $a+b$ is only closable then its closure is again of the form $\widetilde{a} + \widetilde{b}$ where $\widetilde{a}, \widetilde{b}$ are obtained by the usual limiting process and Theorem 4.1 applies. We omit the details.

5 A conjecture on $\|(I+AC)^{-1}\|$

One of our issues was a bound for $\|(I+AC)^{-1}\|$. We conjecture the following, more elegant and possibly tighter bound for this quantity

Conjecture 5.1 *Let A, C be Hermitian positive semidefinite. Then*

$$\|(I+AC)^{-1}\| \leq \|I+AC\| \quad (51)$$

and the equality sign is attained if and only if $AB=0$.

Proof in two dimensions. We obviously may always suppose one of the matrices as diagonal, that is,

$$A = \begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix}$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ \bar{c}_{12} & c_{22} \end{bmatrix}$$

with $a_{11}, a_{22}, c_{11}, c_{22} \geq 0$ and $c_{11}c_{22} \geq |c_{12}|^2$. Now,

$$I+AC = \begin{bmatrix} 1+a_{11}c_{11} & a_{11}c_{12} \\ a_{22}\bar{c}_{12} & 1+a_{22}c_{22} \end{bmatrix}$$

and

$$\begin{aligned} \Delta = \det(I+AC) &= 1 + a_{11}c_{11} + a_{22}c_{22} + a_{11}a_{22}(c_{11}c_{22} - |c_{12}|^2) \\ &\geq 1 + a_{11}c_{11} + a_{22}c_{22} \geq 1. \end{aligned}$$

Hence

$$\begin{aligned} (I + AC)^{-1} &= \frac{1}{\Delta} \begin{bmatrix} 1 + a_{22}c_{22} & -a_{11}c_{12} \\ -a_{22}c_{12} & 1 + a_{11}c_{11} \end{bmatrix} \\ &= \frac{1}{\Delta} U(I + AC)^T U^T \end{aligned} \quad (52)$$

where

$$U = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Thus,

$$\|(I + AC)^{-1}\| = \frac{1}{\Delta} \|I + AC\| \quad (53)$$

and (51) follows. In the case of equality we must have $\Delta = 1$ which implies

$$a_{11}c_{11} = 0, \quad a_{22}c_{22} = 0, \quad a_{11}a_{22}(c_{11}c_{22} - |c_{12}|^2) = 0$$

which, together with the positive semidefiniteness of A, C , implies $AC = 0$. Q.E.D.

Proof for $A = uu^*$, where u is a column vector (A a dyadic) and C arbitrary. This is reduced to the previous case in the following way. As it is well known, there is always a unitary matrix Q such that

$$Qu = \begin{bmatrix} \hat{u} \\ 0 \end{bmatrix}, \quad QCu = \begin{bmatrix} \hat{v} \\ 0 \end{bmatrix},$$

where \hat{u}, \hat{v} are two-dimensional vectors. Now,

$$Q(I + uu^*C)Q^* = \begin{bmatrix} I_2 + \hat{u}\hat{u}^* & 0 \\ 0 & I \end{bmatrix}.$$

On the other hand,

$$Q(I + uu^*C)Q^* = I + Quu^*Q^*QCQ^* = \begin{bmatrix} I_2 + \hat{u}\hat{u}^* & 0 \\ 0 & I \end{bmatrix} QCQ^*.$$

Thus,

$$I_2 + \hat{u}\hat{u}^* = I_2 + \hat{u}\hat{u}^*\hat{C},$$

where \hat{C} is some 2×2 - Hermitian, positive semidefinite matrix, so here the theorem is known to hold. Thus,

$$\|(I + uu^*C)^{-1}\| = \max\{1, \|(I + \hat{u}\hat{u}^*)^{-1}\|\} \leq \max\{1, \|I + \hat{u}\hat{u}^*\|\} = \|I + uu^*C\| = \|I + \hat{u}\hat{u}^*\hat{C}\|,$$

where the last equality follows from (3.2). In the case of equality we have

$$\|(I + \hat{u}\hat{u}^*)^{-1}\| = \|I + \hat{u}\hat{u}^*\|$$

which implies $\hat{u}\hat{u}^* = \hat{u}\hat{u}^*\hat{C} = 0$ that is, $uu^*C = 0$. Q.E.D.

In this case we note the formula

$$(I + AC)^{-1} = I - \frac{AC}{1 + u^*Cu} \quad (54)$$

which may be useful in computations.

Another case in which (51) obviously holds, is when A, C commute. Further support for its general validity is the experimental evidence with randomly generated semidefinite matrices A, C . The observed quotient

$$\frac{\|(I + AC)^{-1}\|}{\|I + AC\|}$$

was always less than one and it approached one only when AC was small.

However, the same numerical experiments have also shown that both (51) and (11) still could be gross overestimates for $\|(I + AC)^{-1}\|$. This, of course, immediately implies that our main estimate (14) is probably not optimal as well.

6 Boundary conditions and invertibility — a case study.

The most prominent example of an operator whose invertibility depends on boundary conditions is the Laplacian on an interval with Dirichlet and Neumann boundary conditions. A deeper manifestation of this phenomenon is encountered in the spectral analysis of Schrödinger operators with periodic potential. Such operators exhibit a spectrum consisting of intervals so called spectral bands. If one restricts the Schrödinger operator originally defined on \mathbb{R} or \mathbb{R}^d to a finite interval or cube, respectively, it is desirable to preserve the periodic structure of the original, unrestricted operator as much as possible. A restriction to a finite cube with Dirichlet boundary conditions leads to spurious eigenvalues located in the spectral gaps of the original operator. A consistent way to avoid these boundary-induced eigenvalues is to impose periodic or, more generally, quasi-periodic boundary conditions. For such restrictions the arising spectrum is contained in the spectrum of the original operator, see [10] for an exposition for operators on $L^2(\mathbb{R})$. In the context of periodic Schrödinger operators, spectral pollution in gaps is a well studied subject, see e. g. [1].

In this section we want to explore these ideas applied to a block-operator investigated in the recent paper [2]. There the following matrix of order $n = 2m$ is considered

$$H_c = H_c(n) = \begin{bmatrix} A + 2cI & B \\ -B & -A - 2cI \end{bmatrix} \quad (55)$$

with the $m \times m$ -blocks

$$A = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 1 & \dots & \vdots \\ 0 & 1 & \ddots & & \vdots \\ \vdots & & & 0 & 1 \\ 0 & \dots & \dots & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ -1 & 0 & 1 & \dots & \vdots \\ 0 & -1 & \ddots & & \vdots \\ \vdots & & & 0 & 1 \\ 0 & \dots & \dots & -1 & 0 \end{bmatrix} \quad (56)$$

where c is any real number (the factor 2 is set by convenience) and $n = 2m$.

We will analyse the spectrum of H_c and see that it exhibits two spurious eigenvalues. To remove these, we will introduce a low-rank modification \tilde{H}_c concentrated on the “boundary”. This results in a circulant-type matrix mimicking periodic boundary conditions used in the context of periodic Schrödinger operators. The specific type of the circulant matrix shows that the operator considered in [2] lives on the two-fold covering space $\{1, \dots, 2m\} \rightarrow \{1, \dots, m\}$.

Of course, the spectrum of $H_c = H_c(n)$ will depend on the dimension n , so we will say that an interval \mathcal{I} around zero is a *global spectral gap* of H_c , if $\mathcal{I} \cap \sigma(H_c) = \emptyset$ for all n and

\mathcal{I} is the largest interval with this property.

We perform the off-diagonalisation by taking the unitary matrix

$$U = \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ I & -I \end{bmatrix} \quad (57)$$

and obtaining

$$K_c = U^{-1}H_cU = UH_cU = \begin{bmatrix} 0 & A_c - B \\ A_c + B & 0 \end{bmatrix} = 2 \begin{bmatrix} 0 & T_c \\ T_c^* & 0 \end{bmatrix}, \quad (58)$$

with

$$T_c = \begin{bmatrix} c & & & & \\ 1 & c & & & \\ & 1 & \ddots & & \\ & & \ddots & c & \\ & & & 1 & c \end{bmatrix} \quad (59)$$

(all void places are zeros). As it is well known, the eigenvalues of K_c , including multiplicities, consist of \pm singular values of T_c . Now, the latter are of some importance in Matrix Numerical Analysis, see [3] where it was shown that the smallest singular value of T_c tends to zero for $m \rightarrow \infty$ and any fixed c with $|c| < 1$. In any case the singular values of T_c are independent of the sign of c as is seen from the property

$$U_0 T_c U_0 = T_{-c}, \quad [U_0]_{ij} = (-1)^j \delta_{ij}.$$

We will now study these singular values in some detail. We shall distinguish the cases

- (i) $c = 0$
- (ii) $0 < c < 1$
- (iii) $c = 1$
- (iv) $c > 1$

For $c = 0$ the matrices T_c, T_c^* are partial isometries with all singular values equal to 1, except for

$$\mathcal{N}(T_c) = \text{span}\{e_m\}, \quad \mathcal{N}(T_c^*) = \text{span}\{e_1\}$$

where e_1, \dots, e_m is the canonical basis in \mathbb{R}^m . Hence K_c has the eigenvalues ± 2 each with multiplicity $n - 1$ and a double eigenvalue zero with

$$\mathcal{N}(K_c) = \text{span} \left\{ \begin{bmatrix} 0 \\ e_m \end{bmatrix}, \begin{bmatrix} e_1 \\ 0 \end{bmatrix} \right\}. \quad (60)$$

The case $c > 1$ allows an easy information based on the representation (55), because then the matrix

$$A_c := A + 2cI = \begin{bmatrix} 2 & 1 & 0 & \dots & 0 \\ 1 & 2 & 1 & \dots & \vdots \\ 0 & 1 & \ddots & & \vdots \\ \vdots & & & 2 & 1 \\ 0 & \dots & \dots & 1 & 2 \end{bmatrix} + (2c - 2)I$$

is positive definite being a sum of two obviously positive definite matrices, so $\sigma(A_c) \geq 2c - 2$. Therefore H_c is quasidefinite and by (2) the interval

$$(-2c + 2, 2c - 2)$$

is contained in the global spectral gap of H_c (and of K_c). For further information we use the fact that the singular values of T_c are square roots of the eigenvalues of $T_c^* T_c$ or, equivalently, of

$$W_c = T_{-c}^* T_{-c} = \begin{bmatrix} c^2 + 1 & -c & 0 & \dots & 0 \\ -c & c^2 + 1 & -c & \dots & \vdots \\ 0 & -c & \ddots & & \vdots \\ \vdots & & & c^2 + 1 & -c \\ 0 & \dots & \dots & -c & c^2 \end{bmatrix}. \quad (61)$$

Now $W_c x = \lambda x$ is componentwise written as

$$\begin{aligned} (c^2 + 1)x_1 - cx_2 &= \lambda x_1 \\ -cx_{j-1} + (c^2 + 1)x_j - cx_{j+1} &= \lambda x_j, \quad j = 2, \dots, m-1 \\ -cx_{m-1} + c^2 x_m &= \lambda x_m \end{aligned}$$

or as a standard second order difference equation

$$-cx_{j-1} + (c^2 + 1 - \lambda)x_j - cx_{j+1} = 0, \quad j = 1, \dots, m \quad (62)$$

with the boundary conditions

$$x_0 = 0, \quad x_m - cx_{m+1} = 0. \quad (63)$$

The solutions

$$x_j = \sin j\alpha, \quad \text{with } \lambda = c^2 + 1 - 2c \cos \alpha \quad (64)$$

and

$$x_j = \sinh j\alpha, \quad \text{with } \lambda = c^2 + 1 - 2c \cosh \alpha \quad (65)$$

automatically satisfy $x_0 = 0$. The second boundary condition from (63) will determine the values of α . This gives

$$(1 - c \cos \alpha) \sin m\alpha - c \cos m\alpha \sin \alpha = 0, \quad (66)$$

$$(1 - c \cosh \alpha) \sinh m\alpha - c \cosh m\alpha \sinh \alpha = 0, \quad (67)$$

respectively. In the easiest case $c = 1$ the substitution (64). immediately leads to

$$\alpha = \alpha_k = \frac{2k-1}{2m+1}\pi, \quad k = 1, \dots, m,$$

giving rise to the eigenvalues

$$\lambda = \lambda_k = 4 \sin^2 \frac{2k-1}{2m+1}\pi, \quad k = 1, \dots, m.$$

In this case the lowest eigenvalue $\lambda_1 = 4 \sin^2 \frac{2}{2m+1}\pi \approx (2m+1)^{-2}$ tends to zero as m tends to infinity, so the global spectral gap of H_c is empty.

In the case $c > 1$ the equation (66) can be written as

$$f(\alpha) = \tan m\alpha \frac{1 - c \cos \alpha}{\sin \alpha} - c = 0, \quad 0 < \alpha < \pi, \quad (68)$$

The localisation of these roots is a bit involved, because there are several different cases to be distinguished. A generic situation is shown on Figure 3 which displays

- The function $f(\alpha)$ (blue) with its poles and roots,
- The function $\lambda = c^2 + 1 - 2c \cos \alpha$ (red), which taken at the roots, gives the eigenvalues,
- the point $\hat{\alpha} = \arccos \frac{1}{c}$ on the α -axis on which f is generically negative.




Figure 3: Functions f and λ for $n = 7$ and $c = 2$

Thus, between each two poles there is a root, except for the two poles enclosing $\hat{\alpha}$; these two poles enclose two roots, altogether n of them. The case in which $\hat{\alpha}$ coincides with one of the poles must be treated separately. But to determine the exact position of the global spectral gap it is enough to notice that in any case for m large enough, the interval

$$0 < \alpha < \hat{\alpha},$$

on which the factor $1 - c \cos \alpha$ is positive, will contain several of the singularities

$$\frac{(2k-1)\pi}{2m}, \quad k = 1, 2, \dots, m$$

of the function f in (68), see Figure 3. Each two of these singularities enclose a root α of (68), and since each of them tends to zero for $m \rightarrow \infty$ the lowest root tends to zero as well.

Hence the corresponding eigenvalue λ from (64) tends to $(c-1)^2$. Since we already know that the interval $(-2(c-1), 2(c-1))$ is contained in the global spectral gap of K_c (and H_c), this interval is, in fact, *equal* to the this gap.

In the case $c < 1$ the factor $1 - c \cos \alpha$ in (68) is globally positive, so the m singularities

$$\frac{(2k-1)\pi}{2m}, k = 1, 2, \dots, m$$

will enclose $m-1$ roots of the equation (68) and the lowest of them will again approach zero with growing m . That is, the corresponding $m-1$ eigenvalues λ from (64) will be larger than $(c-1)^2$ and the lowest of them will approach $(c-1)^2$. Thus, we would have a global spectral gap $(-2(1-c), 2(1-c))$, but for one eigenvalue which is still to be determined. However, as we know from [3] the smallest eigenvalue tends to zero with growing m . This completes the picture. Thus, for $0 \leq c < \infty$ the interval

$$(-2|c-1|, 2|c-1|)$$

is the global spectral gap for K_c (and H_c), **except** that for $0 < c < 1$ there are two 'spurious eigenvalues' tending to zero with growing m .

There is some interest in obtaining the asymptotic estimate of the small eigenvalues. In [3] it was shown that the small singular value of T_c is bounded from above by $\mathcal{O}(\frac{1}{m})$. Numerical experiments indicate that the decay is, in fact, much faster. The setting of difference equations makes it possible to determine the decay accurately and this is what we shall now do.

This solution is obtained by the ansatz $x_j = \sinh j\alpha$ and (67) can be written as

$$g(\alpha) = \tanh m\alpha \frac{1 - c \cosh \alpha}{\sinh \alpha} - c = 0, \quad 0 < \alpha < \infty. \quad (69)$$

From

$$g(0_+) = m(1-c) - c, \quad g(\infty) = -2c$$

by $0 < c < 1$ for m large the equation (69) has a positive root $\alpha = \alpha_1$ which completes the $m-1$ roots previously found whereas the corresponding eigenvalue $\lambda = \lambda_1$ is given by (65). As an approximation to α_1 we propose the value

$$\alpha_0 = \operatorname{arcosh} \frac{c^2 + 1}{2c}.$$

Then a straightforward calculation gives

$$g(\alpha_0) = \frac{-2ce^{-2m\alpha_0}}{1 + e^{-2m\alpha_0}} = -2ce^{-2m\alpha_0} + \mathcal{O}(e^{-2m\alpha_0})$$

and

$$g'(\alpha_0) = \frac{mc}{\cosh^2 m\alpha_0} - \tanh m\alpha_0 \frac{-2c}{1-c^2} = -\frac{2c}{1-c^2} + \mathcal{O}(e^{-2m\alpha_0})$$

Thus, the difference $\delta\alpha = \alpha_1 - \alpha_0$ is given by

$$\delta\alpha = -\frac{g(\alpha_0)}{g'(\alpha_0)} + \mathcal{O}(e^{-2m\alpha_0}) = \mathcal{O}(e^{-2m\alpha_0})$$

whereas the corresponding eigenvalue λ_1 of $W_c = W_c(m)$ is given by

$$\mathcal{O}(e^{-2m\alpha_0}) + \lambda_1 = \delta\alpha(-2 \sinh \alpha_0) = -\frac{4c\delta\alpha}{1-c^2} = 4ce^{-2m\alpha_0}$$

where we have used the fact that the function $c^2 + 1 - 2c \cosh \alpha$ vanishes at $\alpha = \alpha_0$. Hence the small eigenvalues of K_c (and H_c) are asymptotically absolutely bounded by

$$\mathcal{O}(e^{-m\alpha_0}).$$

We summarize the main findings in

Theorem 6.1 *The spectrum of H_c is symmetric w.r.t. zero, i.e. if λ is an eigenvalue, $-\lambda$ is one as well — including multiplicities.*

If $c \geq 1$, the interval $(-2|c-1|, 2|c-1|)$ is a global spectral gap, i.e. $(-2|c-1|, 2|c-1|) \cap \sigma(H_c(m)) = \emptyset$ for all $m \in \mathbb{N}$.

For $c \in [0, 1]$ and each $m \in \mathbb{N}$, $(-2|c-1|, 2|c-1|) \cap \sigma(H_c(m))$ consists of exactly two eigenvalues with absolute value of order $\mathcal{O}(e^{-m\alpha_0})$.

In both cases $(-2|c-1|, 2|c-1|)$ is the maximal interval with the above properties, more precisely for any $\varepsilon > 0, N \in \mathbb{N}$ there exists an $M \in \mathbb{N}$ such that $(-2|c-1| - \varepsilon, 2|c-1| + \varepsilon)$ contains $2N$ eigenvalues of $H_c(m)$ for all $m \geq M$.

The spurious eigenvalues can be computed with high relative accuracy by iteratively solving the equation (69). Under high relative accuracy we mean to obtain a significant number of correct digits *independently* of the size of the computed quantity. Note that the usual matrix computing software computes a singular value of a matrix A with the error of the order $\varepsilon \|A\|$ (ε the machine precision) which may yield no significant digits, if the singular value itself is very small. A notable exception are bidiagonal matrices, which is the case with our T_c . Then there exists an algorithm (and it is implemented in LAPACK and MATLAB packages) which computes each singular value with about the same number of significant digits, no matter how small or how large it may be (barring underflow).²

It is also worthwhile to note that the components $x_j = \sinh j\alpha_1$ of the corresponding eigenvector always agglomerate on one side of the sequence $1, \dots, m$ that is, on the boundary, while all other eigenvectors exhibit standard oscillatory behaviour.

Removing spurious eigenvalues. The form of the null-space of K_0 suggests to introduce the matrix

$$\tilde{K}_c = K_c + \begin{bmatrix} e_m & 0 \\ 0 & e_1 \end{bmatrix} \begin{bmatrix} -2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} e_m^T & 0 \\ 0 & e_1^T \end{bmatrix} = 2 \begin{bmatrix} e_m e_m^T & T_c \\ T_c^* & -e_1 e_1^T \end{bmatrix} \quad (70)$$

For $c = 0$ this leaves all eigenvectors of K_0 unchanged and raises the zero eigenvalues to ± 2 , respectively, thus 'purging' the spurious eigenvalues. The spectrum of \tilde{K}_0 is $\{\pm 2\}$ with the multiplicity m . In particular,

$$\tilde{K}_0^2 = 4I \quad (71)$$

It is a remarkable fact that for $c \neq 0$ the eigenvalues of the matrix \tilde{K}_c still come in plus/minus pairs, including multiplicity. To see this we take

$$J = \begin{bmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 1 & 0 \\ \vdots & & & \vdots \\ 1 & 0 & \cdots & 0 \end{bmatrix} \quad (72)$$

and set

$$\mathbb{P} = \begin{bmatrix} I & 0 \\ 0 & J \end{bmatrix}$$

²In fact, in order to perform with high relative accuracy MATLAB will need the input matrix to be *upper* bidiagonal, so the MATLAB function *svd* should be applied not to T_c but to its transpose.

and note that $Je_1 = e_m$, $Je_m = e_1$ and that the matrix $S_c = T_c J$ is symmetric. Then

$$\mathbb{P}^{-1}\tilde{K}_c\mathbb{P} = \mathbb{P}\tilde{K}_c\mathbb{P} = \begin{bmatrix} e_1 e_1^T & S_c \\ S_c & -e_1 e_1^T \end{bmatrix}.$$

This matrix is of the form (40) and such matrices have the eigenvalues in plus/minus pairs where A and B are allowed to be any symmetric matrices ([9]). It remains to determine the global spectral gap of \tilde{K}_c . In order to do this it is convenient to turn back to the original representation (55) and to form the matrix

$$\tilde{H}_c = U\tilde{K}_c U = \begin{bmatrix} \tilde{A} + 2cI & \tilde{B} \\ \tilde{B}^* & \hat{A} - 2cI \end{bmatrix} \quad (73)$$

with

$$\tilde{A} = \begin{bmatrix} 1 & 1 & 0 & & & \\ 1 & 0 & 1 & & & \\ & 1 & \ddots & \ddots & & \\ & & \ddots & \ddots & \ddots & \\ & & & & 0 & 1 \\ & & & & 1 & -1 \end{bmatrix}, \quad \tilde{B} = \begin{bmatrix} 1 & 1 & 0 & & & \\ -1 & 0 & 1 & & & \\ & -1 & \ddots & \ddots & & \\ & & \ddots & \ddots & \ddots & \\ & & & & 0 & 1 \\ & & & & -1 & 1 \end{bmatrix},$$

$$\hat{A} = \begin{bmatrix} 1 & -1 & 0 & & & \\ -1 & 0 & -1 & & & \\ & -1 & \ddots & \ddots & & \\ & & \ddots & \ddots & \ddots & \\ & & & & \ddots & 0 & -1 \\ & & & & -1 & -1 \end{bmatrix}.$$

By (71) we have $\tilde{H}_c^2 = 4I$ which implies

$$\begin{aligned} \tilde{A}^2 + \tilde{B}\tilde{B}^* &= 4I, \\ \tilde{A}\tilde{B} + \tilde{B}\hat{A} &= 0, \\ \tilde{B}^*\tilde{B} + \hat{A}^2 &= 4I. \end{aligned}$$

Using this³ we see that

$$\tilde{H}_c^2 = \begin{bmatrix} (4 + 4c^2)I + 4c\tilde{A} & 0 \\ 0 & (4 + 4c^2)I - 4c\hat{A} \end{bmatrix} \quad (74)$$

Noting the identity

$$J\hat{A}J = -\hat{A}$$

we see that the eigenvalues of \tilde{H}_c^2 are

$$4 + 4c^2 - 4c\kappa_j, \quad j = 1, \dots, r$$

³Of course, these three identities could be proved directly.

(with the multiplicity two) where κ_j are the eigenvalues of \widehat{A} . These are obtained from the difference equation

$$x_{j-1} + \kappa x_j + x_{j+1} = 0, \quad j = 1, \dots, m \quad (75)$$

with the boundary conditions

$$x_0 = -x_1, \quad x_{m+1} = x_m. \quad (76)$$

The substitution

$$x_j = A \cos j\alpha + B \sin j\alpha$$

solves (75) with

$$\kappa = -2 \cos \alpha$$

whereas the boundary conditions (76) yield after some computation

$$\alpha = \alpha_k = \frac{2k-1}{2m} \pi, \quad k = 1, \dots, m$$

and hence (cf. the Appendix)

$$\kappa = \kappa_k = -2 \cos \frac{2k-1}{2m} \pi, \quad k = 1, \dots, m \quad (77)$$

Thus the eigenvalues of \widetilde{H}_c^2 are

$$4 + 4c^2 + 2c \cos \frac{2k-1}{2m} \pi, \quad k = 1, \dots, m$$

(each taken twice) which is always larger than $4(1-c)^2$ and for m large the set of these eigenvalues comes arbitrarily close to $4(1-c)^2$. Again we conclude:

Theorem 6.2 *The global spectral gap of \widetilde{H}_c is*

$$(-2|c-1|, 2|c-1|).$$

More precisely, for all $m \in \mathbb{N}$ and $c \geq 0$ the eigenvalues of $\widetilde{H}_c(m)$ come in plus/minus pairs and $(-2|c-1|, 2|c-1|) \cap \sigma(\widetilde{H}_c(m)) = \emptyset$. This interval is the largest with this property.

In particular, $(-2|c-1|, 2|c-1|)$ contains no spurious eigenvalues whatsoever.

Finally, we consider the 'infinite dimensional limits' that is, the matrices $\mathbf{H}_c, \mathbf{A}, \mathbf{B}, \mathbf{T}_c, \mathbf{W}_c$, obtained from H, A, B, T_c, W_c by stretching to infinity in both directions. Thus $\mathbf{A} = \mathbf{A}^*$, $\mathbf{B} = -\mathbf{B}^*$, \mathbf{T}_c , and $\mathbf{W}_c = \mathbf{T}_{-c}^* \mathbf{T}_{-c} = \mathbf{W}_c^*$ are bounded operators on the Hilbert space $\ell^2(\mathbb{Z})$, while \mathbf{H}_c and \mathbf{K}_c are selfadjoint operators on $\ell^2(\mathbb{Z}) \oplus \ell^2(\mathbb{Z})$. The operator

$$\mathbf{U} = \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ I & -I \end{bmatrix}$$

is unitary on $\ell^2(\mathbb{Z}) \oplus \ell^2(\mathbb{Z})$, where now I denotes the identity on $\ell^2(\mathbb{Z})$. The operators keep their algebraic relations

$$\mathbf{K}_c = \mathbf{U}^{-1} \mathbf{H}_c \mathbf{U} = \mathbf{U} \mathbf{H}_c \mathbf{U} = \begin{bmatrix} 0 & \mathbf{A}_c - \mathbf{B} \\ \mathbf{A}_c + \mathbf{B} & 0 \end{bmatrix} = 2 \begin{bmatrix} 0 & \mathbf{T}_c \\ \mathbf{T}_c^* & 0 \end{bmatrix}$$

Formula $\mathbf{W}_c = \mathbf{T}_{-c}^* \mathbf{T}_{-c}$ gives us

$$(\mathbf{W}_c x)_j = -c x_{j-1} + (c^2 + 1)x_j - c x_{j+1}$$

Now using the isometric isomorphism $\ell^2(\mathbb{Z}) \rightarrow L^2(0, 2\pi)$

$$\psi(\lambda) = \frac{1}{\sqrt{2}} \sum_{k=-\infty}^{\infty} e^{ij\lambda} x_j$$

the operator \mathbf{W}_c goes over into

$$\begin{aligned} (\mathbf{X}_c\psi)(\lambda) &= \frac{1}{\sqrt{2}} \sum_{j=-\infty}^{\infty} e^{ij\lambda} (-cx_{j-1} + (c^2 + 1)x_j - cx_{j+1}) \\ &= \frac{1}{\sqrt{2}} \sum_{j=-\infty}^{\infty} e^{ij\lambda} (c^2 + 1 - 2c \cos \lambda)x_j = (c^2 + 1 - 2c \cos \lambda)\psi(\lambda) \end{aligned}$$

which is a multiplication operator with the spectrum $[-(1+c)^2, -(1-c)^2] \cup [(1-c)^2, (1+c)^2]$ thus implying the spectral gap $(|1-c|, |1+c|)$ of \mathbf{H}_c . Thus, the obvious approximation H_c , obtained by cutting a 'window' out of \mathbf{H}_c gives rise to spectral pollution in the spectral gap of \mathbf{H}_c . By adding convenient boundary conditions H_c we obtain the modification \tilde{H}_c . This new approximation to \mathbf{H}_c

- (i) has no spectral pollution and
- (ii) keeps the symmetry of the spectrum with respect to zero.

Some numerical experiments. Here we would like to report some interesting observations based on numerical experiments. They are motivated by physical models of disordered systems. In this context the matrix A in (55) is replaced by $V_\omega = A + \text{diag}(\omega_1, \dots, \omega_n)$, where ω_i are independent random variables. Here we will consider a uniform distribution on the interval $[a, b]$. Then, as expected, the multiple eigenvalues ± 2 of the matrix $H_\omega = H \begin{bmatrix} A_\omega & B \\ -B & -A_\omega \end{bmatrix}$ smear into uniformly distributed intervals, but *the double small eigenvalue is only slightly perturbed in the sense that for n large these two eigenvalues tend to zero*. We illustrate this by exhibiting those eigenvalues of the matrix H_ω which are close to zero by taking $a = -3$, $b = 3$.

m = 20	m = 50	m = 100
-7.2091e-01	-3.5965e-01	-0.170365
-5.4659e-01	-1.4649e-01	-0.098804
-5.4215e-04	-4.5522e-10	-9.819153e-31
5.4215e-04	4.5522e-10	9.819153e-31
5.4659e-01	1.4649e-01	0.098804
7.2091e-01	3.5965e-01	0.170365

We emphasize that the exhibited digits of 9.819153e-31 are accurate. The phenomenon of two very small eigenvalues is independent of the choice of a, b . Note that the spurious eigenvalues are not only small but about exponentially small as in the case of the constant diagonal, i.e. $\omega_1 = \dots = \omega_m = 2c$, studied above.

Next we produce a series of graphically represented numerical results with

$$[a, b] = [M - \delta, M + \delta],$$

$n = 100$, $\delta = 0.5$, and M taking the values 0.1, 1, 1.5, 1.8, 2.5, 3. They are contained in Figure 4




Figure 4: Spectral gaps

The upper line (blue) in a pair shows the spectrum of H_ω and the lower (red) the one of \tilde{H}_ω , obtained from H_ω as in (73).

Summarising we may say: *For $M \approx 2$ or so the punctured spectral gap shrinks to zero; then it starts growing again, but small eigenvalues are present no more, because the matrix H_ω has now become quasi-definite.⁴ No theoretical explanation for spurious small eigenvalues in this case seems to be available as yet. On the other hand, as expected, the matrix \tilde{H}_ω lacks the spurious small eigenvalues altogether. With \tilde{H}_ω there is no more symmetry of the spectrum with respect to zero.*

⁴Note that M as the mean value of ω_i corresponds to the value $2c$ with H_c in (55).

Acknowledgement. We are grateful to the colleagues who have obliged us with illuminating discussions and comments. These are W. Kirsch, T. Linß, I. Nakić, M. Omladič, M. Skrzipek, and G. Stolz.

References

- [1] Cancès, E., Ehlacher, V., Maday, Y., Periodic Schrödinger operators with local defects and spectral pollution, *SIAM Journal on Numerical Analysis*, **50** (2012) 3016–3035.
- [2] Chapman, J., Stolz, G., Localization for random block operators related to the XY spin chain, <http://arxiv.org/abs/1308.0708>.
- [3] Erxiong, J., Bounds for the smallest singular value of a Jordan block with an application to eigenvalue perturbation, *LAA* **197** (1994) 697–707.
- [4] Hladnik, M., Omladič, M., Spectrum of the product of operators, *Proc. Amer. Math. Soc.* **102** (1988) 300–302.
- [5] Hong, Y., Horn, R. A., The Jordan Canonical Form of a Product of a Hermitian and a Positive Semidefinite Matrix, *LAA* **147** (1991) 373–386.
- [6] George, A., Ikramov, Kh., A. B. Kucherov, A.B., Some properties of symmetric quasi-definite matrices, *SIAM J. Matrix Anal. Appl.* **21** (2000) 1318–1323.
- [7] Kato, T., *Perturbation Theory for Linear Operators*, Springer 1966.
- [8] van Kempen, H., Variation of the eigenvalues of a special class of Hermitian matrices upon variation of some of its elements, *LAA*, **3** (1970), 263–273.
- [9] Kirsch, W., Metzger, B., Müller, P., Random Block Operators, *J. Stat. Phys.* **143** (2011) 10351054.
- [10] Mezincescu, G. A., Internal Lifschitz singularities for one dimensional Schrödinger operators *Comm. Math. Phys.*, **158** (1993) 315–325.
- [11] Saunders, M. A., Solution of sparse rectangular systems using lsqr and craig, *BIT*, **25** (1995), 588–604.
- [12] Thaller, R., *The Dirac Equation*, Springer 1992.
- [13] Thompson R. C. The Eigenvalues of a Partitioned Hermitian Matrix Involving a Parameter, *LAA* **9** (1974) 243–260.
- [14] Veselić, K., Spectral perturbation bounds for selfadjoint operators I, *Operators and Matrices*, **2** (2008) 307–340.
- [15] Winklmeier, M., *The Angular Part of the Dirac Equation in the Kerr-Newman Metric: Estimates for the Eigenvalues*, Ph. D. Thesis 2005.

Appendix. Some auxiliary computations. Since the matrices of the type (31) appear to be the source of many illustrative examples we give here the explicit formula for their eigenvalues (they come in plus/minus pairs). By setting

$$A = \begin{bmatrix} a_+ & a \\ \bar{a} & a_- \end{bmatrix}, \quad B = \begin{bmatrix} b_+ & b \\ \pm b & b_- \end{bmatrix}$$

such that in the case of the minus sign in B the diagonal elements b_{\pm} are purely imaginary. A straightforward calculation gives

$$(\lambda^2)_{1,2} = \frac{s}{2} \pm \sqrt{s^2 - |a_+ a_- - b_+ b_- - |a|^2 + |b|^2|^2 - |a_+ b_- - b_+ a_- - 2\Re \bar{a} b|^2}. \quad (78)$$

with

$$s = \frac{a_+^2 + a_-^2 + |b_+|^2 + |b_-|^2}{2} + |a|^2 + |b|^2.$$

We derive the formula (77). The substitution $x_j = A \cos j\alpha + B \sin j\alpha$ into (75) gives

$$\begin{aligned} & A \cos j\alpha \cos \alpha + A \sin j\alpha \sin \alpha B \sin j\alpha \cos \alpha - B \cos j\alpha \sin \alpha \\ & + \kappa(A \cos j\alpha + B \sin j\alpha) + \\ & A \cos j\alpha \cos \alpha - A \sin j\alpha \sin \alpha B \sin j\alpha \cos \alpha + B \cos j\alpha \sin \alpha \\ & = 0 \end{aligned}$$

or

$$(A \cos j\alpha + B \sin j\alpha)(\cos \alpha + \cos \alpha + \kappa) = 0$$

thus giving

$$\kappa = -2 \cos \alpha.$$

The boundary conditions (76) give

$$A = -A \cos \alpha - B \sin \alpha$$

$$A \cos(m+1)\alpha + B \sin(m+1)\alpha = A \cos m\alpha + B \sin m\alpha$$

which is a homogeneous linear system

$$\begin{aligned} (1 + \cos \alpha)A + B \sin \alpha &= 0 \\ A(\cos(m+1)\alpha - \cos m\alpha) + B(\sin(m+1)\alpha - \sin m\alpha) &= 0 \end{aligned}$$

so its determinant must vanish:

$$(1 + \cos \alpha)2 \cos \frac{(2m+1)\alpha}{2} \sin \frac{\alpha}{2} + 2 \sin \alpha \sin \frac{(2m+1)\alpha}{2} \sin \frac{\alpha}{2} = 0,$$

equivalently,

$$\begin{aligned} 0 &= \cos \frac{\alpha}{2} \cos \frac{(2m+1)\alpha}{2} + \sin \frac{\alpha}{2} \sin \frac{(2m+1)\alpha}{2} = \\ & \cos \left(\frac{\alpha}{2} - \frac{(2m+1)\alpha}{2} \right) = \cos m\alpha. \end{aligned}$$

Hence

$$\alpha = \alpha_k = \frac{2k+1}{2m} \pi$$

and (77) follows.